



# Pasture Management Recommendations

## GrazonNext® HL Herbicide

### Broadleaf Weed Control with Residual

Broadcast Rate: 1.5–2 pts/A 2 GrazonNext HL<sup>5</sup>

Spot Treatment Rate: 0.25–.5% GrazonNext HL<sup>5</sup>

### Key Points

- The most complete broadleaf weed solution for range & pasture
- Convenient, broad spectrum broadleaf weed control
- Season-long performance with one application
- No livestock grazing restrictions<sup>1</sup>
- Most states require no applicator's license for purchase or application

## PastureGard® HL Herbicide

### Broadleaf Weed Control without Residual

Broadcast Rate: 1–2 pts/A3 PastureGard HL<sup>5</sup>

Spot Treatment Rate: 0.5–1%3 PastureGard HL<sup>5</sup>

### Brush Control (Including Fencerows)

Broadcast Rate: 2–4 pts/A4 PastureGard HL<sup>5</sup>

Spot Treatment Rate: 0.75–1.5%4 PastureGard HL<sup>5</sup>

### Key Points

- Excellent choice where residual herbicides are not wanted
- Controls broadleaf weeds and many woody plants
- No livestock grazing restrictions except lactating dairy animals
- Does not contain 2,4-D
- Ideal for brush control along fence rows and non-irrigation ditch banks
- Premier non-residual weed control when combined with 2, 4-D

## Amount Of Product (In Ounces) to Add to Various Size Tanks

| Sprayer Size | 0.25%   | 0.50%   | 0.75%  | 1%     | 1.50%   |
|--------------|---------|---------|--------|--------|---------|
| 1 Gal        | 0.32 oz | 0.64 oz | 1 oz   | 1.3 oz | 2 oz    |
| 3 Gal        | 1 oz    | 2 oz    | 3 oz   | 4 oz   | 5.75 oz |
| 5 Gal        | 1.5 oz  | 3 oz    | 5 oz   | 6.5 oz | 9.5 oz  |
| 10 Gal       | 3 oz    | 6.5 oz  | 9.5 oz | 13 oz  | 19 oz   |
| 25 Gal       | 8 oz    | 16 oz   | 24 oz  | 32 oz  | 48 oz   |

Always read and follow label directions

1. Restrictions do apply to the transfer of grazing animals to sites intended to grow sensitive crops. Refer to label for a complete list of transfer restrictions
2. For late season applications use the highest recommended rate of 2.1 pts/A GrazonNext HL
3. For broader spectrum non-residual weed control, add 1–2 pts/A 2,4-D for broadcast and 1–2% 2,4-D for spot treatments
4. For broader spectrum control, add 1.5 pts/A GrazonNext HL for broadcast and 0.75% GrazonNext HL for spot treatments
5. It is recommended to add 0.25–0.5% non-ionic surfactant (NIS) for all applications



Label precautions apply to forage treated with GrazonNext HL and to manure from animals that have consumed treated forage within the last three days. Consult the labels for full details. <sup>®</sup>Trademarks of Dow AgroSciences, DuPont or Pioneer and their affiliated companies or respective owners. GrazonNext HL is not registered for sale or use in all states. Contact your state pesticide regulatory agency to determine if a product is registered for sale or use in your state. Always read and follow label directions. ©2019 Corteva Agriscience CR01 000 033 (04/19) 010-58759

Visit us at [rangeandpasture.com](http://rangeandpasture.com)



## GrazonNext® HL

HERBICIDE

## PastureGard® HL

HERBICIDE



### Summary of Restrictions

|                                     | GrazonNext® HL  | PastureGard® HL   |
|-------------------------------------|---|---|
| When Reseeding Grasses              | <p><b>Preemergence:</b> Grasses may be reseeded in the fall following an application of GrazonNext HL applied in the spring or early summer.</p> <p><b>Postemergence:</b> During the season of establishment, GrazonNext HL should be applied only after perennial grasses are well established.</p>  | <p><b>Before Reseeding:</b> Do not reseed treated areas for a minimum of three weeks after treatment.</p> <p><b>Following Reseeding:</b> To avoid grass injury, do not apply until grass seedlings are well established as indicated by tillering (usually after 4 true leaves have emerged), development of a secondary root system and vigorous growth.</p> |
| Plant-Back Restriction              | Do not rotate to cropland for one year following application. Cereals and corn can be planted one year after treatment. Do not plant forage legumes until a soil bioassay has been conducted to show no adverse affect.   | Only wheat, barley, oats or perennial forage grasses may be planted in treated fields within 120 days following application of PastureGard HL.  |
| Grazing or Harvesting               | No haying or grazing restrictions. However, do not transfer animals grazing or feeding on hay to areas where sensitive crops occur without first allowing 3 days of grazing on an untreated pasture.  | <p><b>Lactating Dairy</b><br/>Do not allow grazing or harvest green forage until next growing season. No restrictions for other livestock.</p>  |
| Haying (Harvesting of Dried Forage) | Do not harvest forage for hay within 7 days of application. Cutting hay too soon after spraying weeds can compromise weed control. Wait 14 days prior to cutting grass hay to allow for maximum herbicide activity. Do not move hay made from treated grass within the preceeding 18 months off farm unless allowed by supplemental labeling. | Do not harvest hay within 14 days after application.  |